








Simulation session


Corey Adams (ANL), Makoto Asai (SLAC)

14:00 → 17:00 Day 1 afternoon

Conveners: Corey Adams (ANL) , Makoto Asai (SLAC)

14:00	Simulations: Introduction	🕒 5m
14:05	Event Generation and Simulation Needs for the EIC Speaker: Dr Markus Diefenthaler (Jefferson Lab)  Diefenthaler-AI4EIC...	🕒 25m
14:30	Bottlenecks and limitations in classical simulations: where can AI help? Speaker: Sylvester Joosten (Argonne National Laboratory)  20210907-AI4EIC-S...	🕒 25m
14:55	What AI offers for EIC simulations Speaker: Michelle Kuchera (Davidson College)  Kuchera-AI4EIC-ex...	🕒 30m
15:25	break	🕒 15m
15:40	Generative AI applications for simulations in colliders Speaker: Benjamin Nachman  SimulationAI4EIC_....	🕒 30m
16:10	Machine Learning for the LHCb simulations Speaker: Lucio Anderlini (INFN)  Machine Learning f...	🕒 25m
16:35	Discussion	🕒 25m

Event Generation and Simulation Needs for the EIC



Markus Diefenthaler



The role of AI/ML in simulations

Lesson learned High-precision QCD measurements require high-precision simulations

Statistical accuracy for precise hypothesis testing

- up to trillion of simulated events required (HL-LHC)
- often computationally intensive, in particular calorimeter simulations

Common alternatives

- fast simulations with computationally efficient approximations, e.g., parameterizations or look-up tables
- **still** insufficient accuracy for high-precision measurements

Promising alternatives

- fast generative models, e.g., GANs or VAEs
- AI driven design, e.g., Bayesian optimization

Eur. Phys. J. A (2021) 57:100
https://doi.org/10.1140/epja/s10050-020-00290-x

THE EUROPEAN
PHYSICAL JOURNAL A



Review

A.I. for nuclear physics

Paulo Bedaque¹, Amber Boehnlein^{2,4}, Mario Cromaz³, Markus Diefenthaler², Latifa Elouadrhiri², Tanja Horn⁴, Michelle Kuchera⁵, David Lawrence⁶, Dean Lee⁶, Steven Lidia⁶, Robert McKeown⁷, Wally Melnitchouk², Witold Nazarewicz⁸, Kostas Orginos^{2,7}, Yves Roblin⁹, Michael Scott Smith⁸, Malachi Schram⁹, Xin-Nian Wang³

¹ University of Maryland, College Park, MD, USA

² Thomas Jefferson National Accelerator Facility, Newport News, VA, USA

³ Lawrence Berkeley National Laboratory, Berkeley, CA, USA

⁴ Catholic University, Washington D.C., USA

⁵ Davidson College, Davidson, NC, USA

⁶ Michigan State University, East Lansing, MI, USA

⁷ College of William & Mary, Williamsburg, VA, USA

⁸ Oak Ridge National Laboratory, Oak Ridge, TN, USA

⁹ Pacific Northwest National Laboratory, Richland, WA, USA

Received: 9 September 2020 / Accepted: 7 October 2020 / Published online: 22 March 2021

© Jefferson Science Associates, LLC, under exclusive licence to Società Italiana di Fisica and Springer-Verlag GmbH Germany, part of Springer Nature 2021

Communicated by Ulf Meißner

Abstract This report is an outcome of the workshop AI for Nuclear Physics held at Thomas Jefferson National Accelerator Facility on March 4–6, 2020

Disclaimer: This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DE-AC05-06OR2177. Participation of students and early career professionals was supported by NSF, Division of Physics, under the Grant ‘Artificial Intelligence (AI) Workshop in Nuclear Physics,’ Award Number 2017170. Support for the Hackathon was provided by the University of Virginia School of Data Sciences and by Amazon Web Services.

This report is an outcome of the workshop AI for Nuclear Physics held at Thomas Jefferson National Accelerator Facility on March 4–6, 2020. The workshop brought together 184 scientists to explore opportunities for Nuclear Physics in the area of Artificial Intelligence. The workshop consisted of plenary talks, as well as six working groups.

*e-mail: amber@jlab.org (corresponding author)

Contents

1 Executive summary	2
2 Priority research directions	3
2.1 Future prospects	3
2.2 Community identified needs and commonalities	5
3 Workshop overview	6
4 Summary of workshop sessions	6
4.1 Lattice QCD and other quantum field theories	6
4.1.1 Case studies and future prospects	7
4.1.2 Enabling discoveries/what is needed	7
4.2 Low-energy nuclear theory	7
4.2.1 Current status	7
4.2.2 Case studies and future prospects	8
4.2.3 Enabling discoveries/what is needed	9
4.3 Accelerator science and operations	10
4.3.1 Accelerator and material design optimization	10
4.3.2 Provenance and prognostication for accelerator sub-systems	10
4.3.3 Dynamic optimization of real time operation controls	10
4.3.4 Summary and Final Thoughts	11
4.4 Experimental methods	12
4.4.1 Current status	12
4.4.2 Case Studies and Future Prospects	12
4.4.3 Enabling discoveries/what is needed	13
4.5 Event generation and simulation	13
4.5.1 Current status	13
4.5.2 Case studies and future prospects	14

Springer

Summary

- **Simulations** essential for design of experiments, data analysis, and verification of measurements.
- **Simulations** for the **EIC**, i.e. MCEGs and fast and full detector simulations for the EIC, require **R&D**. We miss core capabilities and we need to work towards accuracy and precision.
- **Simulation R&D** is most efficiently done in common projects and in collaboration with other fields, e.g., HEP or data science.
- Many opportunities for AI/ML to complement and improve **simulations**. While AI/ML approaches will substitute part of **simulation workflows**, they will not replace core tools, e.g., general-purpose MCEGs or Geant4.



BOTTLENECKS IN CLASSICAL SIMULATIONS: WHERE CAN AI HELP?




SYLVESTER JOOSTEN
sjoosten@anl.gov

SIMULATION BOTTLENECKS


Many particles, many components, many steps

- Usually bottlenecks occur where the particle count is high, e.g. as part of a calorimeter shower, or optical photons in a RICH.
- Bottlenecks can also occur in when navigating very detailed geometries (e.g. fiber calorimeters with millions of fibers).
- Finally, scenarios where we need many precise steps through a magnetic field (for upstream & downstream near-beamline detection) is relatively expensive.
- Often multiple bottlenecks at once.



SUMMARY


- Short-term: need large-scale simulations for optimization of a complex detector system.
- Long-term: need (even larger)-scale simulations to properly analyze high-luminosity/high-precision measurements.
- Bottlenecks usually a combination between many particles, many geometry elements and/or many simulation steps.
- Calorimetry, Cherenkov detectors and the far-forward/far-backward regions are prime targets for AI-driven acceleration.



What can AI offer for Simulation at the EIC?




4EIC



Generative Adversarial Networks

Simulation



caloGAN (ATLAS)

Paganini, et. al [1712.10321]

Silicon- Tungsten
calorimeter of the proposed
International Large Detector
Buhmann , et. al
[2005.05334]




Fig. 5 Examples of individual 50 GeV photon showers generated by Geant4 (left), the GAN (center left), WGAN (center right), and BIB-AE (right) architectures. Colors encode the deposited energy per cell.

Summary

- **AI has the potential for large impacts on Simulation for the EIC.**
- **Large body** of prior related work. Often at “bleeding edge” of AI research. Less commonly used for simulation in practice. Requires work.
- **Simulation R&D** is most efficiently done in common projects and in collaboration with other fields, e.g., HEP or data science.
- **Do not expect replacement** of core tools, e.g., general-purpose MCEGs or Geant4.



Generative ML applications for simulations in colliders

Benjamin Nachman

Lawrence Berkeley National Laboratory

bpnachman.com


bpnachman@lbl.gov



@bpnachman



bnachman



AI4EIC


September 7, 2021

Timing

Generation Method	Hardware	Batch Size	milliseconds/shower
GEANT4	CPU	N/A	1772 ←
CALOGAN	CPU <i>Intel Xeon E5-2670</i>	1	13.1
		10	5.11
		128	2.19
		1024	2.03
	GPU <i>NVIDIA K80</i>	1	14.5
		4	3.68
		128	0.021
		512	0.014
		1024	0.012 ←

(clearly these numbers have changed as both technologies have improved - this is simply meant to be qualitative & motivating!)

Integration into real detector sim.



The ATLAS Collaboration fast simulation (AF3) now includes a GAN at intermediate energies for pions

Conclusions and Outlook

42


Generative models hold great promise for enhancing, supplanting, and extending simulations for collider physics

The examples I gave today were not comprehensive

- see the [Living Review](#) for more references

All of the techniques I discussed today could be used for **physics at the EIC!**

This is a link to a recent Berkeley workshop dedicated to generative models →





Istituto Nazionale di Fisica Nucleare
SEZIONE DI FIRENZE




Machine Learning for the LHCb Simulations

Lucio Anderlini on behalf of the LHCb Collaboration
Istituto Nazionale di Fisica Nucleare – Sezione di Firenze


2021 - 09 - 07

Standard simulation: the big picture




Our ML building block

$$\underbrace{x}_{\text{Target variable}} = F^{-1} \left(\underbrace{u}_{\text{Random noise}}; \underbrace{p, \eta, \dots}_{\text{Conditions}} \right)$$




Approaches to ML in simulation




S. Vallecorsa *et al.* EPJ Web Conf., 214 (2019) 02010

Approaches to ML in simulation



F. Ratnikov et al. [EPJ Web Conf., 245 \(2020\) 02026](#)

Approaches to ML in simulation



A. Maevskiy et al. [J. Phys.: Conf. Ser. 1525 012097 \(2020\)](#)

Conclusion 2

Many studies are ongoing to use machine learning to speed up the simulation, taking different approaches:

- simulating the radiation-matter interaction faster
- simulating the response of some detector
- simulating the whole simulation pipeline to reconstructed analysis level quantities

The various solutions are all important as they can all help speeding the simulation up.

To sum up

- Simulation is one of the least advanced areas in the use of AI/ML.
 - Many ideas, opportunities
 - Require works – resource allocation needed
- Two phases
 - Simulation-driven detector optimization with boost of AI/ML (~ 5 years)
 - Fully tailored simulation with maximum use of AI/ML to meet the statistics requirements (~ decade(s))
- Do not expect AI/ML to fully replace core tools, e.g. MCEGs or Geant4.
 - Training, validation
- Modular approach
 - Calorimeter, optical photon detector for PID, very-forward detectors, machine-detector interface, background (neutron, beam halo, etc.) are good start.
- Simulation R&D is most efficiently done in common project and in collaboration with other fields. Engagement with broader AI/Comp.Sci. community is valuable
 - Expect engagement to include data sharing and bidirectional contributions